

Khalid Matrougui, PhD
Bibliography

PUBLICATIONS:

1. Modar Kassan, Maria Galan, Mohamed Trebak, **Khalid Matrougui**. Endoplasmic reticulum stress is involved in aneurysm induction in type 2 diabetes. 2012 Manuscript under writing
2. Sookyong Choi, Modar Kassan, Maria Galan, Mohamed Trebak, **Khalid Matrougui**. NADPH activity impairs vascular function through endoplasmic reticulum stress in type 2 diabetes. 2012 Submitted for publication
3. Modar Kassan, Sookyong Choi, Maria Galan, Alex Bishop, Kazuo Umezawa, Mohamed Trebak, **Khalid Matrougui**. NF κ B Impairs Vascular Function Through PARP-1, SP-1 and COX2-Dependent Mechanisms in Type 2 Diabetes. 2012 Submitted for publication
4. José C. González-Cobos, Xuexin Zhang, Wei Zhang, Brian Ruhle, Rajender K. Motiani, Rainer Schindl, Martin Muik, Jonathan M. Bisailon, Arti V. Shinde, Marc Fahrner, Harold A. Singer, **Khalid Matrougui**, Margarida Barroso, Christoph Romanin & Mohamed Trebak. Store-Independent Orai1/3 channels Activated by Intracrine LeukotrieneC4: Role in Neointimal Hyperplasia in Rats. 2012 Under revision in Journal Clinical Investigation
5. Amin AH, Choi SK, Partyka M, Osman-Elazeik YA, Badr El-Din NK, Kevil C, Navar LG, **Matrougui K**. Sodium Nitrite Therapy Augments Ischemia-Induced Neovascularization and Blood Flow Recovery in Hypertensive Mice. 2011 American Journal of Hypertension (Under Revision)
6. Emerging role of G protein-coupled receptors in microvascular myogenic tone. Gilles Kauffenstein, Ismail Laher, Khalid Matrougui, Nathalie C Guérineau and Daniel Henrion. 2012 Cardiovascular Research – Review (In Press)
7. Kassan M, Galan M, Partyka M, Trebak M, **Matrougui K**. Endoplasmic reticulum stress is involved in cardiac damage and vascular endothelial dysfunction in hypertensive mice. 2012 Arterioscler Thromb Vasc Biol. (In Press)
8. Maria Galan M, Kassan M, Partyka M, **Matrougui K**. Chronic Inhibition of Endoplasmic Reticulum Stress Improves Microvascular Function and Reduces Cardiac Fibrosis in Streptozotocin-Induced Diabetes Mellitus Mice. Hypertension 2012 (In Press)
9. Modar Kassan, Maria Galán, Soo-Kyoung Choi, **Khalid Matrougui**. Endoplasmic Reticulum Stress and Microvascular Endothelial Dysfunction in Diabetes. Editorial - 2012 Journal of Diabetes and Metabolism (In Press)
10. Choi S, Kassan M, Galan M, Partyka M, Trebak M, **Matrougui K** Poly(ADP-ribose) Polymerase-1 (PARP-1) Inhibition Improves Coronary Arteriole Function in Type 2 Diabetes. Hypertension. 2012 May;59(5):1060-8

Khalid Matrougui, PhD
Bibliography

11. Amin A, Choi S, Galan M, Partyka M, Trebak M, Belmadani S, **Matrougui K.** Involvement of Endoplasmic Reticulum Stress and Inflammation in Impaired Ischemia-Induced Neovascularization in Type II Diabetic Mice. *J Pathol. 2012 Jun;227(2):165-74.*
12. Choi S, Galan M, Partyka M, **Matrougui K.** Inhibition of EGFR tyrosine kinase and ERK1/2 MAP-kinase Enhances Ischemia-Induced Neovascularization in type 2 diabetes. *Am J Pathol. 2012 Jan;180(1):410-8.*
13. Zhang W, Halligan KE, Zhang X, Bisaillon JM, Gonzalez-Cobos JC, Hu G, Vincent PA, Zhou J, Singer HA, **Matrougui K** & Trebak M. Orai1-mediated ICRAC is essential for neointima formation after vascular injury. *Circ Res. 2011 Aug 19;109(5):534-42.*
14. Kassan M, Galan M, Partyka M, Trebak M, **Matrougui K.** Interleukin-10 Released by CD4⁺CD25⁺ Natural Regulatory T Cells Improves Microvascular Endothelial Function Through Inhibition of NADPH Oxidase Activity in Hypertensive Mice. *Arterioscler Thromb Vasc Biol. 2011 Nov;31(11):2534-42.*
15. Ali Hassan, Zakaria Abd Elmageed, Megan Partyka, Desiree I Palen, **Khalid Matrougui.** Differential Effects of GRB2-SOS, JNK, STAT and Akt Down Stream Signaling to EGFR Tyrosine Kinase in Mice Coronary Arteriolar Myogenic Tone. *Microvasc Res. 2011 Jan;81(1):135-42*
16. **Matrougui K**, Abd Elmageed Z, Modar K, Choi S, Nair D, Gonzalez Villalobos RA, Chentoufi AA, Belmadani S, Partyka M. Natural Regulatory T Cells and inflammation control Coronary Arteriolar Endothelial Dysfunction in Hypertensive Mice. *Am J Pathol. 2011 Jan;178(1):434-41* (The manuscript was the highlight by the editor in January 2011 Issue)
17. Hans CP, Feng Y, Naura AS, Troxclair D, Zerfaoui M, Siddiqui D, Jihang J, Kim H, Kaye AD, **Matrougui K**, Lazartigues E, Boulares AH. Opposing roles of PARP-1 in MMP-9 and TIMP-2 expression and mast cell degranulation in dyslipidemic dilated cardiomyopathy. *Cardiovasc Pathol. 2010 May 6*
18. Hassan A, Partyka M, Belmadani S, **Matrougui K.** Modified Multipotent Stromal Cells with Epidermal Growth Factor Restore Vasculogenesis and Blood Flow in Ischemic Hind-Limb of Type 2 Diabetic Mice. *Lab Invest. 2010 Jul 90;985-96*
19. Hans CP, Feng Y, Naura AS, Zerfaoui M, Rezk BM, Xia H, Kaye AD, **Matrougui K**, Lazartigues E, Boulares AH. Protective effects of PARP-1 knockout on dyslipidemia-induced autonomic and vascular dysfunction in ApoE mice: effects on eNOS and oxidative stress. *PLoS One. 2009 Oct 13;4(10):e7430.*
20. **Matrougui K.** Diabetes and microvascular pathophysiology: role of epidermal growth factor receptor tyrosine kinase. *Diabetes Metab Res Rev. 2010 Jan;26(1):13-6. Review.*

Khalid Matrougui, PhD
Bibliography

21. Shahid M, Francis J, **Matrougui K**, Majid DS. Involvement of tumor necrosis factor-alpha in natriuretic response to systemic infusion of nitric oxide synthase inhibitor in anesthetized mice. *Am J Physiol Renal Physiol.* 2010 Jul;299(1):F217-24.
22. Shaarawy SM, Tohamy AA, Elgendi SM, Abd Elmageed Z, Bahnasy A, Mohamed MS, **Matrougui K**. Hepatoprotective effects of Garlic and Silymarin in NDEA induced Rats Hepatotoxicity. *Int J Biol Sci.* 2009; 5:549-57
23. Naura A, Hans C, Zerfaoui M, Errami Y, Ju J, Kim H, **Matrougui K**, Kim J, Boulares H. High fat diet induces airway remodeling in ApoE deficient mice: An association with an increase in circulatory and airway inflammatory factors. *Lab Invest.* 2009; 89:1243-51
24. Belmadani S, **Matrougui K**, Kolz C, Palen DI, Prockop D, Chilian W. Amplification of the Atherogenic Aspect of Multipotent Stromal Cells by Epidermal Growth Factor. *Arterioscler Thromb Vasc Biol.* 2009 Jun;29(6):802-8.
25. Hans CP, Feng Y, Naura A, Zerfaoui M, Xia H, Rezk B, **Matrougui K**, E. Lazartigues, Boulares AH. Protective effects of PARP-1 deficiency against dyslipidemia-induced autonomic and vascular dysfunctions in ApoE-/ mice: Role of eNOS activation and oxidative stress. *Plos One.* 2009; 10: e7430
26. Hans CP, Zerfaoui M, Naura A, Troxclair D, Strong JP, **Matrougui K**, Boulares AH. Thieno[2,3-c]isoquinolin-5-one (TIQ-A), a potent poly(ADP-ribose) polymerase inhibitor, promotes atherosclerotic plaque regression in high fat diet-fed ApoE-deficient mice: effects on inflammatory markers and lipid content. *J Pharmacol Exp Ther.* 2009 Apr;329(1):150-8
27. Naura A, Datta R, Hans CP, Zerfaoui M, Rezk B, Errami Y, Oumouna M, **Matrougui K**, Boulares AH. Reciprocal regulation of iNOS and PARP-1 during allergen-induced eosinophilia. *Eur Respir J.* 2009 Feb;33(2):252-62
28. Palen DI and **Matrougui K**. Role of EGFR Phosphorylation in the Induction of Structural Remodeling and Altered Mechanical properties of resistance artery in Type 2 Diabetes. *Diabetes Metab Res Rev* 2008, 651-6
29. Belmadani S, Zerfaoui M, Boulares HA, Palen DI, **Matrougui K**. Role of ERK1/2 MAP-kinase, avb3-integrin shedding and TGFb1 in Resistance Artery Fibrosis in Type-1 Diabetes and Hypertension. *Am J Physio Heart* 2008, H69-76
30. Su J, Lucchesi PA, Suzuki Y, Rezk B, Palen DI, Boulares HA, **Matrougui K**. Role of Advanced Glycation End Products with Oxidative Stress in Resistance Artery Dysfunction in Type 2 Diabetic mice. *Arterioscler Thromb Vasc Biol.* 2008, 1432-8
31. Belmadani S, Palen DI, Gonzalez-Villalobos RA, Boulares HA, **Matrougui K**. Elevated Epidermal Growth Factor Receptor Phosphorylation Induces Resistance Artery Dysfunction in Type 2 Diabetes. *Diabetes* 2008, 57:1-9

Khalid Matrougui, PhD
Bibliography

32. Su J, Palen DI, Boulares HA, **Matrougui K.** Role of ACE-AT1R and ACE-AT2R complexes in the control of mouse mesenteric resistance artery tone. *Molecular and Cellular Biochemistry* 2008 Apr;311(1-2):1-7
33. Ouhtit A, **Matrougui K**, Bengrine A, Koochekpour S, Zerfaoui M, Yousef Z. Survivin is Not only a Death Encounter but a Survival Protein for Invading Tumor Cells. *Front Biosci. 2007;12:1260-70.*
34. Su J, Palen DI, Lucchesi PA, **Matrougui K.** Mice lacking the gene encoding for MMP-9 and resistance artery reactivity. *Biochem Biophys Res Commun. 2006;349(4):1177-8*
35. Palen DI, Ouhtit A, Belmadani S, Lucchesi PA, **Matrougui K.** Hydrogen Peroxide acts as Relaxing Factor in Human Vascular Smooth Muscle Cells Independent of MAP-Kinase and Nitric Oxide. *Front Biosci. 2006;11:2526-34.*
36. Palen DI, Belmadani S, Lucchesi PA, and **Matrougui K.** Role of SHP-1, Kv.1.2 and cGMP in nitric oxide-induced ERK1/2 MAP-Kinase dephosphorylation in rat VSMC. *Cardiovasc Res 2005;268-77.* This article was selected in "The Phosphatase Resource", (www.phosphataseresource.com)
37. Lucchesi PA, Belmadani S, **Matrougui K.** Hydrogen peroxide acts as both vasodilator and vasoconstrictor in the control of perfused mouse mesenteric resistance arteries. *J of Hypertension 2005;571-579.*
38. Lucchesi PA, Belmadani S, Sabri A, **Matrougui K.** Involvement of EGF-receptor trans-activation in myogenic tone induced by pressure but not in contraction induced by angiotensin II in mice mesenteric resistance arteries. *Circulation 2004;110(23):3587-93.*
39. Bergaya S, **Matrougui K**, Meneton P, Henrion D, Boulanger CM. Role of tissue kallikrein in response to flow in mouse resistance arteries. *J of Hypertension 2004 ;22(4):745-50.*
40. Widdop RE, **Matrougui K**, Levy BI, Henrion D. AT2 receptor-mediated relaxation is preserved after long-term AT1 receptor blockade. *Hypertension 2002;40:516-520.*
41. Manlio B, **Matrougui K**, Loufrani L, Maclouf J, Levy B, Levy-Toledano S, Habib A, Henrion D. Thromboxane stimulation in pressurized and perfused resistance arteries. Role of ERK1/2 and p38 MAP-Kinase. *Journal Vascular Research 2002;39:353-360.*
42. Tankó LB, **Matrougui K.** Can We Apply Results From Large To Small Arteries? *Circ Res. 2002;90:68.*
43. Loufrani L, **Matrougui K** Li Z, Levy BI, Lacolley P, Paulin D, Henrion D. Selective Microvascular Dysfunction in Mice Lacking the Gene Encoding for Desmin. *FASEB J 2002;16:117-9.*

Khalid Matrougui, PhD
Bibliography

44. **Matrougui K**, Loufrani L, Levy BI, Henrion D. High salt intake results in inducible of cyclooxygenase 2 and related modifications of flow-induced dilation and pressure-induced myogenic tone in normotensive rats. *Pharmacology & Toxicology* **2001;89:183-187.**
45. **Matrougui K**, Tanko LB, Loufrani L, Gorny D, Levy BI, Tedgui A, Henrion D. Involvement of Rhokinase and actin filament in Mitogen Activated Protein Kinases activity induce by pressure and angiotensin II in intact and pressurised resistance arteries. *Arterioscler Thromb Vasc Biol.* **2001;21:1288-1293.**
46. Tankò LB, Simonsen U, **Matrougui K**, Gregersen H, Frobert O, Bagger JP, Mikkelsen EO. Axial stretch is a modulator of the sensitivity to spasmogen stimuli in porcine large coronary arteries. *Pharmacology & Toxicology* **2001;88:89-97.**
47. Loufrani L, **Matrougui K**, Gorny D, Duriez M, Blanc I, Levy BI, Henrion D. Flow (shear stress)-induced endothelium-dependent dilation is defectuous in mice lacking the gene encoding for dystrophin. *Circulation* **2000;103:864-870.**
48. **Matrougui K**, Eskildsen-Helmond YE, Fiebeler A, Henrion D, Levy BI, Tedgui A, Mulvany MJ. Angiotensin II stimulates Extracellular Signal-Regulated Kinases activity in intact pressurized rat mesenteric resistance arteries. *Hypertension* **2000;36:617-621.**
49. **Matrougui K**, Levy BI, Henrion D. Tissue angiotensin II and endothelin1 modulate differently the response to flow in mesenteric resistance arteries of normotensive and spontaneously hypertensive rats. *Br J Pharmacol* **2000;130(3):521-526.**
50. **Matrougui K**, Loufrani L, Heymes C, Levy BI, Henrion D. Activation of AT(2) Receptors by Endogenous Angiotensin II Is Involved in Flow-Induced Dilation in Rat Resistance Arteries. *Hypertension* **1999;34:659-665.**
51. Iglarz M, **Matrougui K**, Levy BI, Henrion D. Chronic blockade of endothelin ETA receptors improves flow dependent dilation in resistance arteries of hypertensive rats. *Cardiovasc Res* **1998;39:657-64.**
52. **Matrougui K**, Levy BI, Schiavi P, Guez D, Henrion D. Indapamide improves flow-induced dilation in hypertensive rats with a high salt intake. *J of Hypertension* **1998;16:1485-90.**
53. **Matrougui**, Schiavi P, Guez D, Henrion D.. High sodium intake decreases pressure-induced (myogenic) tone and flow-induced dilation in resistance arteries from hypertensive rats. *Hypertension* **1998;32:176-9.**
54. Veksler VI, Lechene P, **Matrougui K**, Ventura-Clapier R. Rigor tension in single skinned rat cardiac cell: role of myofibrillar creatine kinase. *Cardiovasc Res* **1997;36:354-62.**

Khalid Matrougui, PhD
Bibliography

55. Henrion D, Terzi F, **Matrougui K**, Duriez M, Boulanger CM, Colucci-Guyon E, Babinet C, Briand P, Friedlander G, Poitevin P, Levy BI. Impaired flow-induced dilation in mesenteric resistance arteries from mice lacking vimentin. *J Clin Invest* 1997;100:2909-14.
56. **Matrougui K**, Maclouf J, Levy BI, Henrion D. Impaired nitric oxide and prostaglandin-mediated responses to flow in resistance arteries of hypertensive rats. *Hypertension* 1997;30:942-7.
57. Qiu HY, Bernard C, **Matrougui K**, Tedgui A, Levy BI. Rapid effect of LPS-activated macrophages on the reactivity of the rat mesenteric artery. *J Vasc Res* 1996;33:370-9.

SELECTED POSTER AND ORAL COMMUNICATIONS

1. D138 821.3 Interleukin-10 released by natural regulatory T cells improves microvascular endothelium-dependent relaxation in hypertensive mice. M. Kassan, M. Partyka and **K. Matrougui**. Tulane Univ. **EB 2012 Washington**
2. D68 1025.9 PARP-1 inhibition improves coronary arteriole function in type 2 diabetic mice. S-K. Choi, M. Kassan, M. Galan, M. Partyka and **K. Matrougui**. Tulane Univ. **EB 2012 Washington**
3. D162 1030.7 Inhibition of nitric oxide synthase enhances the production of tumor necrosis factor-alpha in macrophage cells. M.T. Islam, D. Agarwal, J. Francis, **K. Matrougui** and D.S. A. Majid. Tulane Univ. Hlth. Sci. Ctr. and LSU Sch. of Vet. Med. **EB 2012 Washington**
4. Inhibition of epidermal growth factor receptor tyrosine kinase and ERK1/2 MAP-kinase enhances ischemia- induced neovascularization in type 2 diabetic mice. S-K. Choi, M. Galan, M. Partyka and **K. Matrougui**. Tulane Univ. (1092.4) **EB 2012 Washington (Oral Presentation and Poster)**
5. A117 1092.3 Reticulum endoplasmic stress and inflammation impair ischemia-induced neovascularization in type 2 diabetic mice. **K. Matrougui**, M. Partyka and A. Amin. Tulane Univ. and Mansoura Univ., Egypt. **EB 2012 Washington**
6. Ali H Amin, Zakaria Y Abd Elmageed, Megan Partyka, Mohammed F Ismail, Osman YA Elazeik, Souad Belmadani, **Khalid Matrougui**. The Role of Adhesion and Migration of Stimulated Multipotent Stromal Cell by EGF in Vasculogenesis in Type 2 Diabetes (**Oral Presentation, AHA 2009 Orlando, Florida**)
7. Zakaria T Abd Elmageed, Devika Nair, Romer A Gonzalez-Villalobos, Aziz A Chentoufi, Megan Partyka, Souad Belmadani, **Khalid Matrougui**, Natural Regulatory T Cells control Coronary Arteriolar Endothelial Dysfunction and Hypertension in Mice (**Poster Presentation, AHA 2009 Orlando, Florida**)

Khalid Matrougui, PhD
Bibliography

8. Zakaria Y Abd Elmageed, Ali H Amin, Devika Nair, Partyka Megan, Souad Belmadani, **Khalid Matrougui**. The Inhibition of Poly(ADP-ribose) Polymerase-1 (PARP-1) Significantly Accelerates the Vasculogenesis in a Model of Hind Limb Ischemia (**Poster Presentation, AHA 2009 Orlando, Florida**)
9. Zakaria Y Abd Elmageed, Ali H Amin, Megan Partyka, Desiree I Palen, **Khalid Matrougui**. Mechanisms of Coronary Arteriolar Myogenic Tone: Role of Down Stream Signaling to EGFR Tyrosine Kinase (**Poster Presentation, AHA 2009 Orlando, Florida**)
10. Zakaria Abd Elmageed, Ali Hassan, Megan Partyka, Desiree I Palen, **Khalid Matrougui**. Differential Effects of GRB2-SOS, JNK, STAT and Akt Down Stream Signaling to EGFR Tyrosine Kinase in Mice Coronary Arteriolar Myogenic Tone. Experimental Biology Meeting, New Orleans, Louisiana 2009
11. Su J, Lucchesi PA, Gonzalez R, Rezk B, Suzuki Y, Palen DI, Boulares HA, **Khalid Matrougui**. Role of Advanced Glycation End Products with Oxidative Stress in Resistance Artery Dysfunction in Type 2 Diabetic mice. Experimental Biology Meeting, San Diego, California 2008
12. Desiree I Palen, **Khalid Matrougui**. Elevated Epidermal Growth Factor Receptor Phosphorylation Induces Resistance Artery Dysfunction in Type 2 Diabetes. AHA meeting 2007, Florida, USA
13. Desiree PI, **Matrougui K**. Eutrophic structural remodeling of hypertensive and type 1 diabetic mouse mesenteric resistance artery (EB-2007, Washington Convention Center, Washington, DC)
14. Desiree PI, **Matrougui K**. Involvement of EGF receptor and oxidative stress in type 2 diabetic mouse resistance artery dysfunction (EB-2007, Washington Convention Center, Washington, DC)
15. Desiree PI, **Matrougui K**. Role of TGF β 1 and avb3-integrin in the control of collagen type 1 turnover in response to Ang II and high glucose (EB-2007, Washington Convention Center, Washington, DC)
16. Desiree PI, **Matrougui K**. Eutrophic structural remodeling of mesenteric resistance artery in Type 2 diabetic mouse (EB-2007, Washington Convention Center, Washington, DC)
17. Su J, Belmadani S, Palen D, Lucchesi P, Boulares H, **Matrougui K**. Mouse Resistance Artery Tone Regulation: Role of the Enzyme Converting Angiotensin and Angiotensin Type 1 - 2 Receptors. AHA Dallas November 13-17, 2005. Poster 1050/B105. Circulation

Khalid Matrougui, PhD
Bibliography

18. Su J, Lucchesi PA, Palen DI, **Matrougui K**. Mice Lacking Gene Encoding for MMP9 Show High Sensitivity to Shear Stress and Alteration of MMP2/9 Involvement in Myogenic Tone in Resistance Artery. *Vascular Biology and Medicine* 2005, Chicago, IL USA
19. Lucchesi PA, Belmadani S, Palen DI, **Matrougui K**. Evidence for Hydrogen Peroxide as Relaxing Factor in Human Vascular Smooth Muscle Cells Independently of MAP-Kinase and Nitric Oxide. *Vascular Biology and Medicine* 2005, Chicago, IL USA
20. Lucchesi PA, Belmadani S, Sabri A, **Matrougui K**, Involvement of MMPs 2-9 in EGF Receptor Transactivation in Pressure-Induced Myogenic Tone in Mouse Mesenteric Resistance Arteries. *Experimental Biology* 2004 annual meeting. Washington. DC, USA. Abstract/Oral # 2315.
21. Lucchesi PA, Belmadani S, **Matrougui K**, Involvement of SHP-1 Tyrosine Phosphatase in Nitric Oxide Induced MAP-Kinase Dephosphorylation in Rat Vascular Smooth Muscle Cells. *Experimental Biology* 2004 annual meeting. Washington. DC, USA. Abstract # 2339.
22. **Matrougui K** et al. Evidence for Hydrogen Peroxide as Endothelium Derived Hypolarization/Relaxing Factor in Human Vascular Smooth Muscle Cells Independently of MAP-Kinase and NO-Synthesis. *Experimental Biology* 2004 annual meeting. Washington. DC, USA. Abstract # 2344
23. **Matrougui K** et al. Evidence for Hydrogen Peroxide as Endothelium Derived Hypolarization/Relaxing Factor in Human Vascular Smooth Muscle cells independently of MAP-Kinase and NO-synthase. *FASEB J* July 5-10, 2003, Denver Colorado.
24. **Matrougui K** et al. Angiotensin II stimulates extracellular signal-regulated kinases activity in intact pressurized rat mesenteric resistance arteries (B121). *J. Submicrosc. Cytol. Pathol.*, 2000; 32: 427. (XIth International Vascular Biology Meeting, Geneva, Switzerland).
25. **Matrougui K**, et al. Involvement of Rho-kinase and the actin filament network in angiotensin II-induced contraction and extracellular signal regulated kinase (ERK1/2) activation in intact and pressurized rat resistance arteries. *Circulation*, 2000; 102; II-314 (1539) (American Heart Association, 2000, 73th Scientific Sessions, New-Orleans, LA, USA).
26. Loufrani L, **Matrougui K**, et al. Flow (shear stress)-induced arterial endothelium-dependent dilation and vascular density are decreased in mice genetically deficient in dystrophin. *Circulation*, 2000; 102; II-224 (1090) (American Heart Association, 2000, 73th Scientific Sessions, New-Orleans, LA, USA).

Khalid Matrougui, PhD
Bibliography

27. Loufrani L, **Matrougui K**, et al. The absence of dystrophin is associated with a decreased flow-induced dilation and a lower arteriolar density in mice. *Hypertension*, 2000; 36:656 (5.06) (European conference on Blood Pressure and Cardiovascular disease, Noordwijkerhout, Pays-Bas, Oct. 2000).
28. Loufrani L, **Matrougui K** et al. Attenuation of Flow-Induced Dilation in Carotid Artery from Mice Lacking Dystrophin Gene. *Arch Mal Coeur Vaiss*, 1999; (Journées d'Hypertension, Paris, 1999).
29. Loufrani L, **Matrougui K**, et al. Attenuation of Flow-Induced Dilation in the Carotid Arteries from Mice Lacking the Gene for Dystrophin. *Circulation*, 1999; 100 (suppl. I): I-415 (abstract. 2181, (American Heart Association, 1999, 72 Scientific Sessions, Atlanta, GA, USA)).
30. **Matrougui K**, et al. Tissue angiotensin II and endothelin-1 antagonize the response to flow in mesenteric resistance arteries of spontaneously hypertensive rats. *Hypertension*, 1999; 34:715 (European conference on Cardiovascular disease, Noordwijkerhout, Pays-Bas, Oct. 1999).
31. **Matrougui K**, et al. Activation of AT₂ receptors by endogenous angiotensin II is involved in flow-induced dilation in rat resistance arteries. *Hypertension*, 1999; 34: 698 (4.02) (European research conference on Blood Pressure and Cardiovascular disease, Noordwijkerhout, Pays-Bas, Oct. 1999).
32. Loufrani L, **Matrougui K**, et al. Attenuation of Flow-Induced Dilation in the Carotid Arteries from Mice Lacking the Gene for Dystrophin. *Hypertension*, 1999; 34: 695 (1.07) (European research conference on Blood Pressure and Cardiovascular disease, Noordwijkerhout, Pays-Bas, Oct. 1999).
33. **Matrougui K**, et al. Attenuation of flow-induced dilation in resistance arteries from mice lacking dystrophin. *Arch Mal Coeur Vaiss*, 1999; 92: 490 (9-12), (Congrès GRRC 1999).
34. **Matrougui K**, et al. High NaCl intake decreases flow-dilation and myogenic tone in resistance arteries from normotensive rats: Role of cyclooxygenase-2. *Arch Mal Coeur Vaiss*, 1999; 92: 494 (9-13), (Congrès GRRC : Biologie et Pathologie du Coeur et des Vaisseaux, Deauville, 1999).
35. Henrion D, Terzi F, **Matrougui K**, et al. Flow-induced dilation in mesenteric resistance arteries from mice lacking vimentin. (3d International Workshop, Versailles, 1998).
36. **Matrougui K**, et al. Decrease in flow-induced dilation and myogenic tone in resistance artery in hypertensive rats fed with high salt: *FASEB J*, 1998; 12; 4:532 (EB, 1998, San Francisco, USA).

Khalid Matrougui, PhD
Bibliography

37. **Matrougui K**, et al. Flow-induced dilation in rat resistance arteries: Role of angiotensin II type 2. Arch Mal Coeur Vaiss, 1998; 91: 488, (Congrès GRRC, Lille, 1998).
38. **Matrougui K**, et al. Involvement of angiotensin II type 2 and endothelin-1 type A receptor stimulation in response to shear stress in rat resistance arteries *in situ*. J. Vasc. Res., 1998; 35 (S1): O22 (5th International Symposium on Resistance Arteries, Meren, Belgique, 1998).
39. **Matrougui K**, et al. Vasodilatation AT2-dépendante et vasoconstriction AT1-dépendante induite par le débit des artères de résistance. Arch Mal Coeur Vaiss, 1997 (Journées d'Hypertension, Paris, 1997).
40. **Matrougui K**, et al. Modification de la vasodilatation AT2-dépendante et de la vasoconstriction AT1-dépendante induite par le débit dans des artères de résistance de rats spontanément hypertendus. Arch Mal Coeur Vaiss, 1997; 90 (Suppl): 11-8, (Congrès GRRC: Arcachon, 1997).
41. **Matrougui K**, et al. Involvement of angiotensin II type 2 and endothelin-1 type A receptor stimulation in response to shear stress in rat resistance arteries *in situ*. Circulation, 1997; 96 (Suppl I): 2671, (American Heart Association, 1997, 70 Scientific Sessions, New Orleans, LA, USA).
42. **Matrougui K**, et al. Prostaglandins-mediated response to flow and prostaglandins production are impaired in hypertensive rat resistance mesenteric arteries. Hypertension, 1996; 28: 41 (50th Scientific Sessions of the Council for High Blood Pressure Research, 1996, Chicago, MI, USA).
43. **Matrougui K**, et al. Flow-induced dilation in resistance arteries *in situ* involves prostaglandins but not nitric oxide (NO) in hypertensive rats. FASEB J, 1996; 10: 4041 (EB, 1996, Washington, DC, USA).